


PROFLAVANOL™ 90



- Supplies a potent combination of grape-seed bioflavonoids and **POLY C™**
- When combined with a healthy diet, helps support sound cardiovascular function, immune function, and circulation*
- Provides advanced antioxidant support*



ITEM# 132

It is an incongruity of life. We require oxygen for efficient energy production and for the function of many metabolic pathways. And yet oxygen, in a variety of chemical forms collectively known as “reactive oxygen species” (ROS), is damaging to our cells and tissues.^{1,2} Elevated exposure to ROS creates a physiological imbalance known as oxidative stress, which is also caused by free radicals.^{3,4}

Free radicals and ROS have become true enemies to our health. Every day these reactive molecules—from environmental pollution, cigarette smoke, fatty diets, and normal cellular metabolism—assault our cells.³⁻⁷ Food supplements rich in antioxidants, such as **PROFLAVANOL™ 90**, support the body’s natural defenses against free radicals.^{8-13*}

PROANTHOCYANIDINS

Proanthocyanidins are one of the most potent antioxidants, and they strongly inhibit ROS. Research suggests a strong role for dietary bioflavonoids in supporting and maintaining sound cardiovascular function.¹⁴⁻¹⁶ This originates from the observation that relatively low rates of heart disease exist in France, despite a diet high in saturated fat.¹⁷ This apparent “French paradox” has been attributed to the proanthocyanidins in the red wine consumed with most meals. Human clinical research has also shown that proanthocyanidins in red wine maintain healthy endothelial function.^{18,19*}

Proanthocyanidins have been studied extensively in Europe and in the United States. Scientifically observed and documented benefits include the following:

- Helps protect plasma LDL from oxidation^{20*}
- Helps retain healthy capillary strength and vascular function^{21,22*}
- Supports healthy immune function^{23*}
- Increased peripheral circulation^{24*}

WHY PROFLAVANOL™ 90?

With USANA’s unique formulation, **PROFLAVANOL 90** combines the highest quality grape-seed extract with the free-radical quenching power of exclusive **POLY C™**, giving you an important tool in the quest for lifelong good health.^{25,26} Laboratory studies done at USANA confirm that, when used together, these ingredients provide significantly better protection against the oxidative modification of low density lipoproteins (LDL) than would be predicted from the sum of their individual effects. Not only do proanthocyanidins neutralise free radicals, they also conserve and regenerate vitamins C and E.²⁷ **PROFLAVANOL 90** offers superior antioxidant support.*

OPTIMIZERS/CARDIOVASCULAR

DIRECTIONS: TAKE ONE (1) TO THREE (3) TABLETS DAILY, PREFERABLY WITH MEALS. DO NOT EXCEED STATED RECOMMENDED DAILY DOSE.

EACH TABLET CONTAINS:	%EC RDA*	
VITAMIN C	300mg	500%
GRAPESEED EXTRACT (VITIS VINIFERA)	90mg	†

*RDA = Recommended Daily Allowance
†No EC RDA Established

INGREDIENTS: VITAMIN C [AS POLY C™ (CALCIUM ASCORBATE, POTASSIUM ASCORBATE, MAGNESIUM ASCORBATE, ZINC ASCORBATE)], BULKING AGENT (MICROCRYSTALLINE CELLULOSE), GRAPESEED EXTRACT, BULKING AGENT (SODIUM CARBOXYMETHYLCELLULOSE), STARCH, ANTI-CAKING AGENT (ASCORBYL PALMITATE), GLAZING AGENT (DEXTRIN), ANTI-CAKING AGENT (SILICON DIOXIDE), GLAZING AGENTS (DEXTRINE, SOYA LECITHIN), ACIDITY REGULATOR (SODIUM CITRATE).

ALLERGY ADVICE: CONTAINS SOYA.

KEEP OUT OF REACH OF YOUNG CHILDREN. CONSULT YOUR DOCTOR IF YOU ARE PREGNANT, LACTATING, TAKING A PRESCRIPTION MEDICATION, OR HAVE A KNOWN MEDICAL CONDITION. FOOD SUPPLEMENTS SHOULD NOT BE USED AS A SUBSTITUTE FOR A VARIED DIET.

References

1. Bland JS. J Nutr Environ Med 1995;5:255-80.
2. Halliwell B. Ann Rev Nutr 1996;16:33-50.
3. Gaziano JM. Proc Assoc Am Physicians 1999;111:2-9.
4. McKeown N. Nutr Rev 1999;57:321-24.
5. Fletcher RH, Fairfield KM. JAMA 2002;287(23):3127-9.
6. Ames BN. Mutat Res 2001;475(1-2):7-20.
7. Ames BN, et al. Proc Natl Acad Sci USA 1993;90(17):7915-22.
8. Frei B ed. Natural Antioxidants in Human Health and Disease. San Francisco: Academic Press; 1994.
9. de Rijk M, et al. Arch Neurol 1997;54:762-65.
10. Papas AM. Antioxidant Status, Diet, Nutrition, and Health. Boca Raton: CRC Press; 1999.
11. Liu S, et al. Am J Clin Nutr 2000;72:922-28.
12. Michaud DS, et al. A J Clin Nutr 2000;72:990-97.
13. Slater TF. Royal Soc Med International Congress and Symp Series 1981;47:11-15.
14. Dell'Agli M, et al. Cardiovasc Res 2004;63(4):593-602.
15. Folts JD. Adv Exp Med Biol 2002;505:95-111.
16. Cos P, et al. Curr Med Chem 2004;11(10):1345-59.
17. Renaud S, de Lorgeril M. Lancet 1992;339:1523-26.
18. Stoclet JC, et al. J Physiol Pharmacol 1999;50: 535-40.
19. Cuevas AM, et al. Lipids 2000;35:143-48.
20. Fuhrman B, et al. Am J Clin Nutr 1995;61:549-54.
21. Gazave JM, et al. Biochem Exp Biol 1978;14:25-31.
22. Baruch J. Ann Chir Plast Esthet 1984;29(4).
23. Kakegawa . et al. Chem Pharm Bull (Tokyo) 1985;5079-82.
24. Beylot C, Bioulac P. Gaz Med de France 1980;87(22).
25. Maffei F, et al. Arzneimittelforschung 1994;44:592-601.
26. Whitehead TP, et al. Clin Chem 1995;41:32-35.
27. Lagrue G, et al. Sem Hop Paris 1981;September:18-25.

***These statements have not been evaluated by the Food Standards Agency. This product is not intended to diagnose, treat, cure, or prevent any disease.**